

METABOLISM-U BLUEPRINT SERIES

BP-033

BUILDING YOUR METABOLICALLY HEALTHY DIET

How to Build a Nutritionally Complete, Calorie-Balanced Way of Eating

Adequacy • Balance • Quality • Individualization • Sustainability

Rodney C. Curtis, MD
Metabolism-U LLC

Metabolism-U Insight
There is no single perfect diet. But there are well-supported principles for constructing a dietary pattern that provides adequate nutrition, supports healthy body composition and metabolic health, and can be sustained over time.

MEDICAL & NUTRITION DISCLAIMER. This Blueprint is provided for general educational and informational purposes only. It is not a substitute for individualized medical care, diagnosis, treatment, or nutrition counseling. Nutritional needs and the safety or appropriateness of dietary changes vary with age, health conditions, medications, kidney and gastrointestinal function, nutritional status, activity, and other individual factors. Before making significant changes to your diet, calorie intake, protein intake, supplements, or other nutrition practices, consider discussing the issues presented here with your personal physician and/or a licensed or appropriately credentialed nutrition professional who can evaluate your individual circumstances. Medical and nutrition evidence, professional guidelines, and recommendations change over time; although this material is reviewed for accuracy at publication, portions may become outdated as new evidence emerges. Do not delay or disregard professional medical advice because of information in this Blueprint.

© 2026 Metabolism-U LLC. All Rights Reserved.

AI USE DISCLOSURE. Artificial intelligence tools were used to assist with research support, literature organization, editing, formatting, and publication preparation. AI-generated or AI-assisted material can contain errors or omissions and does not replace professional medical or nutritional judgment. The content was reviewed and edited for Metabolism-U publication.

Why This Blueprint Matters

Nutrition has become unnecessarily confusing. Low carbohydrate, low fat, ketogenic. Mediterranean plant-based, intermittent fasting, count every calorie, don't count calories at all. Someone trying simply to eat well can understandably wonder: What am I actually supposed to eat?

The fundamentals are less controversial than the noise suggests. Healthy dietary patterns can take many forms, but they repeatedly share several characteristics: appropriate energy intake, adequate protein, plentiful vegetables and fruits, fiber-rich carbohydrate sources, predominantly unsaturated fats, sufficient vitamins and minerals, and moderation of dietary components that become problematic when consumed excessively. Current international guidance similarly emphasizes adequacy, balance, moderation and diversity.[1]

Metabolism-U Insight

The goal of this Blueprint is not to prescribe the perfect diet. It is to teach you how to construct yours.

Step 1 — Define What Your Diet Needs to Accomplish

Before deciding how much protein, carbohydrate and fat to eat, ask a more fundamental question: What am I trying to accomplish? Maintenance, fat loss, body recomposition, muscle gain, athletic performance and prevention of frailty each require different energy and nutrient strategies. Age, body size and composition, activity, medical conditions, medications, nutritional status, culture and preferences also matter. The diet should begin with the person—not with a predetermined macronutrient percentage.

Step 2 — Determine How Much Energy You Actually Need

Calories are not the entire story of nutrition, but energy intake has to match the goal. A useful starting point is to estimate resting energy expenditure (RMR) and then approximate total daily energy expenditure (TDEE). The Mifflin–St Jeor equation is one commonly used predictive equation for adults:[2]

Men: RMR ≈ (10 × weight kg) + (6.25 × height cm) − (5 × age) + 5

Women: RMR ≈ (10 × weight kg) + (6.25 × height cm) − (5 × age) − 161

After estimating RMR, multiply it by an activity factor to obtain an initial estimate of total daily energy expenditure (TDEE). Use the table below as a practical starting guide.

Activity level	Practical description	Starting multiplier
Sedentary	Mostly seated lifestyle; little purposeful exercise; relatively low daily steps.	× 1.2
Lightly active	Regular daily movement plus light exercise about 1–3 days/week; some walking, but largely sedentary occupation.	× 1.35–1.4
Moderately active	Purposeful exercise about 3–5 days/week and/or substantial routine walking and daily movement.	× 1.5–1.6
Very active	Hard exercise most days, high daily movement, physically demanding work, or substantial endurance/resistance training.	× 1.7–1.8
Exceptionally active	Heavy physical work combined with substantial training, high-volume endurance training, or comparable activity.	≈ 1.9+

Activity Example

If calculated resting metabolic rate, RMR, is 1,600 kcal/day: sedentary ≈ 1,920 kcal/day (1,600 × 1.2); lightly active ≈ 2,240 (×1.4); moderately active ≈ 2,480 (×1.55); very active ≈ 2,800 (×1.75). The same person can therefore have substantially different calorie requirements depending on daily movement and exercise.

Activity categories are estimates, not precise measurements. Two people who both exercise four days per week may expend very different amounts of energy if one sits most of the day while the other routinely walks 12,000 steps. Exercise duration and intensity also matter. Use the multiplier to establish a starting estimate, then calibrate it using actual weight and body-composition trends.

Let Your Body Calibrate the Estimate

Suppose an equation predicts maintenance at 2,400 kcal/day. Eat approximately that amount consistently while monitoring average body weight under reasonably standardized conditions (preferably as soon as you wake up and naked). If the weight trend is stable, the estimate is probably close. If weight consistently falls, intake is likely below expenditure; if it rises, intake is likely above expenditure. Use trends, over seven to ten days, rather than single weigh-ins because water, glycogen, sodium, activity level and gastrointestinal contents can move scale weight substantially.

Metabolism-U Insight

The equation gives you an estimate. Your longitudinal response helps calibrate it.

Match Calories to the Goal

Goal	Starting strategy	Example if TDEE ≈ 2,400
Maintain	Start near estimated TDEE	≈2,400 kcal/day
Lose body fat	Moderate deficit	≈1,900–2,100 kcal/day
Recomposition	Near maintenance or modest deficit	≈2,150–2,350 kcal/day
Gain muscle/weight	Modest surplus	≈2,550–2,700 kcal/day

These are illustrative starting ranges—not universal prescriptions. During fat loss, faster is not automatically better, particularly when preservation of lean mass is a priority. During muscle gain, a modest surplus can support training and tissue growth without unnecessarily accelerating fat gain. Recomposition means attempting to reduce fat mass while maintaining or increasing lean mass; scale weight may change little even while waist circumference, strength and body composition improve.

As body weight, activity and training change, energy needs can change too. Reassess rather than assuming the original calorie target remains correct.

Metabolism-U Insight
ESTIMATE → IMPLEMENT → MEASURE → ADJUST

Step 3 — Establish the Protein Foundation

Protein provides the amino acids required to maintain and repair skeletal muscle and other tissues. This becomes particularly important with aging because muscle contributes to strength, mobility, glucose disposal, resilience and physiological reserve.

The adult RDA is 0.8 g/kg/day (about 0.36 g/lb/day), but this is not necessarily an optimal target for an older adult deliberately trying to preserve or build muscle. Expert groups focused on healthy aging commonly recommend about 1.0–1.2 g/kg/day for healthy older adults, with higher intakes often appropriate for physically active adults and during illness or recovery.[3,4] Sports-nutrition guidance commonly places exercising adults around 1.4–2.0 g/kg/day.[5]

For a healthy older adult engaged in progressive resistance training and deliberately trying to maximize muscle preservation or growth, protein intake approaching 1 gram per pound of body weight per day (≈2.2 g/kg/day) may be a reasonable individualized higher-end target. It should not be presented as a universal requirement or as proven superior to somewhat lower adequate intakes. A large resistance-training meta-analysis found that additional gains in fat-free mass plateaued on average around 1.6 g/kg/day, although individual requirements and goals vary.[6] Higher-protein diets warrant individualized guidance when kidney disease or other relevant medical conditions are present.

Protein Quality and the Calorie Budget

High-quality proteins provide all essential amino acids in adequate proportions and are readily digestible. Dairy, eggs, fish, poultry and meat are examples; soy is an important high-quality plant protein. A varied plant-based diet can also provide adequate essential amino acids, but the amount and mixture of foods may need more deliberate planning. Essential amino acids—particularly leucine—are important components of the muscle protein synthetic response.

Protein source also affects the calorie budget. Lean animal proteins and concentrated sources such as whey isolate can provide substantial protein with relatively few accompanying calories. Some whole-food plant protein sources provide protein together with carbohydrate and/or fat, so obtaining the same protein dose can sometimes require more total food and calories. That is not a nutritional disadvantage by itself: beans, nuts, seeds and other plant foods may simultaneously provide fiber, unsaturated fats and micronutrients. The practical question is how the protein source fits the person's total calorie and nutrient requirements.

Metabolism-U Insight
Protein needs are not identical for everyone. Age, activity, calorie intake, training, health and body-composition goals should determine the target.

Step 4 — Build the Nutrient-Rich Core Around Your Needs

Once calorie and protein targets are established, fill the remaining calorie budget deliberately. A metabolically healthy diet should derive much of its energy from vegetables, fruits, legumes, whole grains, nuts, seeds, appropriate protein sources and predominantly unsaturated fats. These foods collectively supply fiber, vitamins, minerals, essential fatty acids and other nutrients.

Adjust the pattern to the problem you are trying to solve. An endurance athlete may allocate more calories to carbohydrate. Someone resistance training may prioritize protein. During fat loss, lean protein sources and high-volume, nutrient-rich foods can make a calorie deficit easier to sustain. During muscle or weight gain, larger portions and energy-dense nutritious foods can help. An older adult with poor appetite may need more protein and nutrient density in less food volume.

Metabolism-U Insight
Don't simply choose healthy foods. Choose amounts and combinations of healthy foods that match the nutritional problem you are trying to solve.

Step 5 — Understand Your Carbohydrates and Fats

Carbohydrates: Faster and Slower Are Context, Not Morality

Carbohydrates range from simple sugars to starches and fiber. Their metabolic effects depend not only on the carbohydrate itself but also on fiber, food structure, processing, accompanying protein and fat, portion size and the meal as a whole.

Highly refined grains, sugary beverages and some low-fiber starches can be digested and absorbed relatively rapidly, producing a faster rise in blood glucose and insulin. That is not automatically harmful; rapidly available carbohydrate can be useful around prolonged or demanding exercise. Problems are more likely when large quantities of rapidly consumed, refined carbohydrates repeatedly add calories while contributing little fiber or nutritional value.

Whole grains, legumes, many vegetables and intact or minimally processed carbohydrate foods often contain more fiber and retain more physical structure, which can slow digestion and glucose appearance and may improve satiety. Glycemic index describes the blood-glucose response to a standardized amount of available carbohydrate; glycemic load also accounts for the amount of carbohydrate in a usual portion. Neither measure alone determines whether a food is healthy.[7]

Metabolism-U Insight

Don't classify carbohydrates simply as good or bad. Ask what kind of carbohydrate it is, what comes with it, how much you are eating, and what you need it for.

Dietary Fat: We Need It, but Type Matters

Fat provides essential fatty acids, participates in cell structure and signaling, and enables absorption of vitamins A, D, E and K. Monounsaturated fats are abundant in olive oil, avocado and many nuts. Polyunsaturated fats include essential omega-6 and omega-3 fatty acids; fatty fish provides EPA and DHA, while nuts, seeds and plant oils provide other unsaturated fatty acids.

Saturated fats occur in fatty meats, butter, full-fat dairy, coconut and palm oils, and many processed foods. Saturated fat need not be described as a poison, but replacing some saturated fat—particularly with polyunsaturated fat—can reduce LDL cholesterol and cardiovascular risk. Industrially produced trans fats should be minimized. WHO recommends that unsaturated fats predominate, with saturated fat below 10% of energy and trans fat below 1%.[1,8]

Fat provides about 9 kcal/g compared with about 4 kcal/g for protein and carbohydrate, so portions matter when calorie control is the goal. A nutritious handful of nuts or tablespoon of olive oil still contributes to the energy budget.

Step 6 — Make Fiber and Micronutrient Adequacy Deliberate

Fiber is not a single substance. Soluble or viscous fibers—found in foods such as oats, barley, legumes, fruit and psyllium—can influence post-meal glucose, satiety and LDL cholesterol; many fermentable fibers also feed intestinal microorganisms. Insoluble fibers, abundant in foods such as wheat bran, whole grains, vegetables, nuts and seeds, add stool bulk and support normal intestinal transit. Many foods contain both types.[9]

For most people, the practical goal is not to calculate separate soluble and insoluble targets but to eat a variety of fiber-rich plant foods. WHO recommends at least 25 g/day of naturally occurring fiber for adults and at least 400 g/day of fruits and vegetables.[1] A calorie-balanced diet can still be deficient in vitamins and minerals, so dietary variety matters. Supplements can appropriately fill identifiable gaps or medical needs, but they should not be expected to rescue an otherwise poorly constructed diet.

Step 7 — Control the Excesses Without Creating Forbidden Foods

A metabolically healthy diet does not require nutritional perfection. What matters most is the overall pattern. Habitual excess calories, free or added sugars, sodium, saturated and trans fats, alcohol, and highly energy-dense foods that are easy to overconsume deserve attention. Current WHO guidance recommends free sugars below 10% of total energy, saturated fat below 10%, trans fat below 1%, and sodium below 2 g/day for the general adult population.[1] These are population-health targets—not evidence that a food suddenly becomes dangerous when a numerical threshold is crossed.

Ultra-processed foods should not dominate the diet, but processing alone does not determine nutritional value. The goal is moderation, not food fear.

Step 8 — Build Meals, Not Just Nutrient Targets

Most people do not eat macronutrient ratios; they eat meals. After estimating calorie and protein requirements, translate those targets into food. A practical meal can begin with protein, abundant vegetables and/or fruit, an appropriate portion of quality carbohydrate, and a source of predominantly unsaturated fat. Then adjust portions to the person's energy needs, protein needs, activity, body-composition goal and medical circumstances.

MF-033-02 — BUILD A METABOLICALLY HEALTHY MEAL

A practical plate framework



Step 9 — Make the Diet Sustainable

A nutritionally sophisticated diet that works only under perfect circumstances is not sustainable. Real life includes restaurants, birthdays, holidays, vacations, travel, family gatherings and meals eaten simply for enjoyment. Build flexibility into the structure: keep most meals anchored by appropriate calories, adequate protein, vegetables and/or fruit, fiber-rich foods and sensible portions, while allowing selected meals or occasions to be more flexible.

When a special meal is planned, modestly simplify other meals if useful; lean protein, vegetables, fruit and lighter portions, without fasting all day or trying to 'earn' food with excessive exercise. At restaurants, identify the protein and vegetables first, choose carbohydrate and fat portions that fit your goal, and decide which extras—bread, appetizers, alcohol or dessert—matter most to you. You do not have to consume every extra simply because it is available, and an occasional celebration does not need to become a calorie-counting exercise.

Holidays and vacations are periods to manage, not tests to pass or fail. Maintenance may be a perfectly reasonable temporary goal. Keep a few anchors in place—protein, some fruits and vegetables, physical activity when practical, reasonable portions and attention to hunger and fullness—then return to your usual structure when the event or trip ends. Do not wait for Monday, next month or January 1.

Expect the scale to move. Restaurant meals, travel, higher sodium or carbohydrate intake, alcohol, changes in training, constipation and the amount of food in the gastrointestinal tract can temporarily change body weight. Glycogen is also stored with water. A rapid increase after a weekend or vacation therefore does not necessarily represent an equivalent gain in body fat. Judge progress by the longer-term trend rather than a single weigh-in.

A temporary departure is data,not failure. There is no need to punish an indulgent meal with starvation or excessive exercise. Resume your normal plan at the next reasonable opportunity. The important distinction is between a temporary departure and allowing that departure to become the new pattern. The sooner you return to your structure, the less important the temporary deviation becomes.

PLAN → ENJOY → ACCEPT NORMAL FLUCTUATIONS → RETURN TO STRUCTURE → CONTINUE

Success is not never departing from the plan. Success is developing a dietary structure you can return to again and again.

The Metabolism-U Diet Construction Framework

DEFINE THE GOAL → ESTIMATE ENERGY REQUIREMENTS → MATCH CALORIES TO THE GOAL → ESTABLISH PROTEIN → BUILD THE NUTRIENT-RICH CORE → CHOOSE QUALITY CARBOHYDRATES & FATS → CHECK FIBER & MICRONUTRIENTS → MODERATE THE EXCESSES → MAKE IT ENJOYABLE & SUSTAINABLE

Metabolism-U Insight

A good diet is not something you pass or fail. It is a nutritional system you measure, adjust and improve.

Continue Building Your Diet at Metabolism-U. For additional practical nutrition guidance, visit metabolismu.com and open the Nutrition & Recipes section, including the Metabolism-U nutrition guidelines and related resources for putting these principles into practice.

Final Thoughts

Nutrition does not need to be a contest between carbohydrates and fat, plants and meat, fasting and frequent meals, or "clean" and "bad" foods. The more useful question is whether your overall diet provides the right amount of energy and enough of the nutrients your body needs while supporting the health, body composition, physical function and life you want.

There is no universally perfect diet. But there is a scientifically sound process for building a very good one.

For more insights into nutrition, consider checking out the Matabololism-U website's Nutrition and Recipe section at www.MetabolismU.com .

Better Metabolism – Stronger Body – More Independent You.

ABOVE ALL — NEVER SETTLE.

Selected References

1. World Health Organization. Healthy diet. Updated January 26, 2026. Accessed September 13, 2026.
2. Mifflin MD, St Jeor ST, Hill LA, Scott BJ, Daugherty SA, Koh YO. A new predictive equation for resting energy expenditure in healthy individuals. Am J Clin Nutr. 1990;51(2):241-247. doi:10.1093/ajcn/51.2.241.
3. Bauer J, Biolo G, Cederholm T, et al. Evidence-based recommendations for optimal dietary protein intake in older people: a position paper from the PROT-AGE Study Group. J Am Med Dir Assoc. 2013;14(8):542-559. doi:10.1016/j.jamda.2013.05.021.
4. Deutz NEP, Bauer JM, Barazzoni R, et al. Protein intake and exercise for optimal muscle function with aging: recommendations from the ESPEN Expert Group. Clin Nutr. 2014;33(6):929-936. doi:10.1016/j.clnu.2014.04.007.
5. Jäger R, Kerksick CM, Campbell BI, et al. International Society of Sports Nutrition position stand: protein and exercise. J Int Soc Sports Nutr. 2017;14:20. doi:10.1186/s12970-017-0177-8.
6. Morton RW, Murphy KT, McKellar SR, et al. A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. Br J Sports Med. 2018;52(6):376-384. doi:10.1136/bjsports-2017-097608.
7. World Health Organization. Carbohydrate intake for adults and children: WHO guideline. Geneva: World Health Organization; 2023.
8. World Health Organization. Saturated fatty acid and trans-fatty acid intake for adults and children: WHO guideline. Geneva: World Health Organization; 2023.
9. Reynolds A, Mann J, Cummings J, Winter N, Mete E, Te Morenga L. Carbohydrate quality and human health: a series of systematic reviews and meta-analyses. Lancet. 2019;393(10170):434-445. doi:10.1016/S0140-6736(18)31809-9.
10. Food and Agriculture Organization of the United Nations; World Health Organization. What are healthy diets? Joint statement. Geneva: WHO; 2024. ISBN 978-92-4-010187-6.

© 2026 Metabolism-U LLC. Complete, unaltered sharing for educational purposes is permitted with attribution.